

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Abraxane	Oncology- Injectable	IV	No	No	Yes	Store at room temp. Protect from bright light. Freezing or refrigerating do not adversely affect the stability of intact vials. Different standards apply after reconstitution	No	Yes
Actemra	Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials/syringes at 2°C to 8°C (36°F to 46°F). Do Not freeze. Protect vials and syringes from light (store in the original package until time of use); keep syringes dry. Solutions diluted for IV infusion in NS may be stored in refrigerator or room temperature for up to 24 hrs; solutions diluted for IV infusion in 1/2NS may be stored at 2°C to 8°C for up to 24 hrs or at room temperature for up to 4 hrs; protect from light. Discard unused product remaining in the vials. Refrigerate. Avoid excessive or vigorous agitation; do Not shake. Discard if intact vial is left at room temperature for >12 hours prior to use. Refrigerate; do Not freeze. Discard unused portions; do Not save for further use. Do Not use if previously frozen	Yes	No
Actimmune	Infectious Disease	Injection SubQ	Yes	Yes	N/A	Limited Distribution product distributed by CVS Specialty. Store intact vials refrigerated at 2°C to 8°C in the original carton. Protect from light. Reconstituted solution should be diluted immediately in NS, D5W, or LR; however, may be stored refrigerated for up to 24 hours; do not freeze. Solutions diluted for infusion should be used immediately after preparation; however, may be stored for 24 hours refrigerated; use within 24 hours of initial reconstitution.	No	No
Adagen	Immune Deficiencies & Related Disorders	Injection IM	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials refrigerated at 2°C to 8°C in the original carton. Protect from light. Reconstituted solution should be diluted immediately in NS, D5W, or LR; however, may be stored refrigerated for up to 24 hours; do not freeze. Solutions diluted for infusion should be used immediately after preparation; however, may be stored for 24 hours refrigerated; use within 24 hours of initial reconstitution.	No	No
Adcetris	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Adcirca	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	No	Yes
Adempas	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Advate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution product distributed by CVS Specialty	Yes	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Adynovate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution product distributed by CVS Specialty	Yes	No
Afinitor	Oncology- Oral/Topical	Oral	Yes	No	Yes	Protect from light; protect from moisture. Prior to reconstitution, store refrigerated; do not freeze. Do not refrigerate after reconstitution. Can also store at room temperature up to 3 months; do not return to refrigerator. Store in original package to protect from light. Use within 4 hours of reconstitution.	No	Yes
Afstyla	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution product distributed by CVS Specialty. Refrigerate; do not freeze. Protect from light. Do not shake. Following dilution in NS, solution for infusion should be used immediately; however, if not used immediately, diluted solutions should be refrigerated at 2°C to 8°C for up to 36 hours.	Yes	No
Aldurazyme	Lysosomal Storage Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store in original container to protect from light and moisture.	Yes	No
Alecensa	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate. do Not freeze or shake.	Yes	No
Alferon N	Infectious Disease	Injection Intralesional	No	No	N/A		Yes	No
Alimta	Oncology- Injectable	IV	No	No	Yes	Reconstituted solution and infusion solution are stable for 24 hours when refrigerated at 2°C to 8°C Products are stable for three years, up to the expiration date printed on the label. For single use; discard any unused contents in vial. Store intact vials at ≤25°C; do not freeze. May store reconstituted solution at room temperature. Use within 3 hours of reconstitution; do not refrigerate after reconstitution.	No	Yes
Alphanate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No		Yes	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Alphanine SD	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	When stored at refrigerator temperature, 2°C to 8°C, factor IX is stable for the period indicated by the expiration date on its label. Avoid freezing which may damage container for the diluent. May also be stored at room temperature not to exceed 30°C for up to 1 month. Reconstituted solution should be used within 3 hours of preparation. Refrigerate. do not freeze. May be stored at room temperature (not to exceed 30°C) for up to 6 months. After removal from refrigeration, do not return to refrigerator. Store in the original package to protect from light. Following reconstitution, may be stored at room temperature (not to exceed 30°C) for no longer than 3 hours. Protect from direct sunlight. Do not refrigerate after reconstitution.	Yes	Yes
Alprolix Alunbrig	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders Oncology- Oral/Topical	IV Oral	No Yes	No No	No Yes	none	Yes No	No No
Ampyra	Multiple Sclerosis	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Apokyn	Movement Disorders	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Aralast NP	Alpha-1 Antitrypsin Deficiency	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Aranesp	Anemia	Injection SubQ or IV	Yes	No	Yes	Store at 2°C to 8°C (36°F to 46°F); do Not freeze. Do Not shake. Protect from light. Store in original carton until use. The following stability information has also been reported: May be stored at room temperature for up to 7 days	No	Yes
Arcalyst	Cryopyrin-Associated Periodic Syndrome	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Arzerra	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Protect from light. Solutions diluted in NS for infusion must be started within 12 hours of preparation if not used immediately); discard any remaining solution 24 hours after preparation.	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Aubagio	Multiple Sclerosis	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Austedo	Movement Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Avastin	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate in original carton; do not freeze. Protect from light; do not shake. Solutions diluted in NS are stable for up to 8 hours under refrigeration. Discard unused portion of vial.	Yes	No
Aveed	Hormonal Therapies	Injection IM	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 25°C; excursions are permitted between 15°C and 30°C. Store Aveed in original container. Refrigerate, Do not freeze. Protect from light. Vial: If refrigeration is not available, may be stored at 25°C (77°F) for up to 30 days	Yes	No
Avonex	Multiple Sclerosis	Injection IM	Yes	Yes	Yes	Prior to reconstitution, store intact vials at room temperature of 25°C; excursions permitted to 15°C to 30°C	No	No
Azacitidine (Vidaza)	Oncology- Injectable	Injection SubQ or IV	Yes	Yes	Yes		No	No
Bebulin	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store undiluted vials at 2°C to 8°C; do not freeze. Limited Distribution product distributed by CVS Specialty. Retain in original package until use. The reconstituted solution may be stored for 12 hours at 15°C to 25°C. Solutions diluted for infusion in NS may be stored for up to 36 hours (including infusion time) at 15°C to 25°C	Yes	Yes
Beleodaq	Oncology- Injectable	IV	No	No	Yes		Yes	No
Bendeka	Oncology- Injectable	IV	No	No	Yes	Store when stored at 15°C to Refrigerate. Avoid freezing which may damage the diluent syringe. Reconstituted solution should be at room temperature and used within 3 hours of preparation. Do not refrigerate after reconstitution.	Yes	No
BeneFIX	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No		Yes	Yes
Benlysta	Systemic Lupus Erythematosus	Injection SubQ or IV	No	No	Yes	Refrigerate. Protect from light. Dispense in original container if applicable	No	Yes
Berninert (Canada Brand Name)	Hereditary Angiodema	IV	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Betaseron	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	If not used immediately following reconstitution, refrigerate solution at 2°C to 8°C and use within 3 hours; do not freeze or shake solution.	No	Yes
Bethkis	Cystic Fibrosis	Nebulized solution	Yes	No	No	Limited Distribution product distributed by CVS Specialty. Store under refrigeration. May be stored in foil pouch (opened or unopened) at room temperature of 25°C for up to 28 days. Protect from light. The colorless to pale yellow solution may darken over time if Not stored under refrigeration; however, the color change does Not affect product quality. Do Not use if solution has been stored at room temperature for >28 days. Store at 2°C to 25°C (36°F to 77°F). Protect from light. Avoid humidity and high temperatures after opening bottle	Yes	No
Bexarotene (Targretin)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Dilution is not recommended. Don't freeze or shake	No	Yes
Bivigam	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Refrigerate. Dilution is not recommended. Don't freeze or shake	Yes	No
Blincyto	Oncology- Injectable	IV	No	No	Yes	ervative free) are stable in NS	Yes	No
Bosulif	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Botox	Batten Disease	Injection IM	No	No	No	Store uNopened vials under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 36 months or until the expiration date on the vial. After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) and use within 24 hours (does Not contain a preservative). Botox 100-unit vial: The following stability information has also been reported: May be stored at room temperature for up to 5 day	No	Yes
Bravelle	Infertility	Injection SubQ or IM	Yes	Yes	No	Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Brineura	Batten Disease	Injection Intraventricular	No	No	Yes	Store cerliponase alfa injection and intraventricular electrolytes injection upright in a freezer at -25°C to -15°C in original carton and protect from light. Store administration kit in original carton separately from cerliponase alfa. Do Not freeze kit. Thaw cerliponase alfa and intraventricular electrolytes vials at room temperature for approximately 60 minutes. Preferably, administer immediately after thawing. May store thawed vials at 2°C to 8°C for 24 hours. Use product held in labeled syringes immediately. If Not used immediately, store in syringes at 2°C to 8°C and use within 4 hours. Vials are single use only. Limited distribution products Not distributed by CVS Specialty	Yes	No
Cabometyx Capecitabine (Xeloda)	Oncology- Oral/Topical Oncology- Oral/Topical	Oral Oral	Yes Yes	No No	Yes Yes	Limited Distribution product distributed by CVS Specialty	Yes No	No Yes
Caprelsa	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	No
Carbaglu	Urea Cycle Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed Not by CVS Specialty. Prior to reconstitution, store at or below 30°C (86°F). Do not freeze. Following reconstitution in a sterile laminar air flow environment, store under refrigeration. Don't freeze, shake, or heat	Yes	No
Carimune NF	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Prior to reconstitution, store at 2°C to 8°C (36°F to 46°F). Once removed from refrigeration, aztreonam and the diluent may be stored at room temperature (up to 25°C [77°F]) for ≤28 days. Protect from light. Use immediately after reconstitution.	No	Yes
Cayston	Cystic Fibrosis	Nebulized solution	Yes	No	N/A	Limited Distribution product distributed by CVS Specialty	Yes	No
Ceprotin	Coagulation Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Cerdelga	Lysosomal Storage Disorders	Oral	Yes	No	Yes	Refrigerate	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Cerezyme	Lysosomal Storage Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate intact vials. Reconstituted solution is stable for 12 hours refrigerated. Solution diluted for infusion in NS is stable for up to 24 hours when refrigerated. Discard unused solution.	Yes	Yes
Cetrotide	Infertility	Injection SubQ	Yes	Yes	No	Lyophilized powder may be stored in the refrigerator or at room temperature of 3°C to 25°C. Protect from light. Discard unused portion.	No	Yes
Cholbam	Gastrointestinal Disorders- Other	Oral	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Chorionic gonadotropin (Novarel, Pregnyl)	Infertility	Injection IM	Yes	Yes	No	Store at intact vials at 15°C to 30°C. Following reconstitution, solution is stable when refrigerated for 30 days (Novarel) or 60 days (Pregnyl). Store intact vials and syringes at 2°C to 8°C; do Not freeze. Do Not separate contents of carton prior to use. Protect from light. Bring to room temperature prior to administration. Store reconstituted solution for up to 24 hours at 2°C to 8°C (do Not freeze); Do Not leave at room temperature for more than 2 hours prior to administration.	No	Yes
Cimzia	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty, Refrigerate	No	Yes
Cinqair	Allergic Asthma	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes
Cinryze	Hereditary Angiodema	IV	Yes	Yes	Yes	Store intact vials at 2°C to 30°C; do not freeze. Protect from light. After reconstitution, use within 1 hr	Yes	No
Coagadex	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution product Not distributed by CVS Specialty	Yes	No
Cometriq	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 2°C to 8°C; do not freeze. Protect from light. May be stored at room temperature for up to 6 mon; do not return to refrigerator if stored at room temp	Yes	No
Corifact	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No		Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Cosentyx	Psoriasis	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. After reconstitution of the vial, use the solution immediately or store in the refrigerator at 2°C to 8°C for up to 24 hours	Yes	Yes
Cotellic	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty Store at 2°C to 8°C (36°F to 46°F) for up to 36 months or at room temperature for up to 12 months; do Not return vial to refrigerator after it has been stored at room temperature. Do Not freeze. Keep in original carton to protect from light.	Yes	No
Cuvitru	Transplant	Injection SubQ or IM	No	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Store intact vials at 2°C to 8°C; do not freeze. Retain in original carton to protect from light. Do not shake. Solutions diluted in NS for infusion may be stored at 2°C to 8°C for no longer than 24 hours (do not freeze) or may be stored for 4 hours at room temperature, do not shake	No	No
Cyramza	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Protect from light and moisture.	Yes	No
Cystagon	Lysosomal Storage Disorders	Oral	Yes	No	No	Limited Distribution product Not distributed by CVS Specialty. Store in freezer. Remove from freezer 24 hours before use; store thawed product for up to 1 week. Discard product 1 week after thawing; do not refreeze.	Yes	No
Cystaran	Lysosomal Storage Disorders	Ophthalmic Solution	Yes	No	No	Refrigerate. Infusion should begin within 6 hours after entering the vial.	Yes	No
Cytogam	Immune Deficiencies & Related Disorders	IV	No	No	Yes	No	Yes	No
Daklinza	Hepatitis C	Oral	Yes	No	Yes	No	No	Yes
Darzalex	Oncology- Injectable	IV	No	No	Yes	ation may be kept at room t	Yes	No



Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Decitabine (Dacogen)	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Vials should be reconstituted with 10 mL SWFI to a concentration of 5 mg/mL. Immediately further dilute with NS or D5W to a final concentration of 0.1 to 1 mg/mL. Use appropriate precautions for handling and disposal. Solutions not administered within 15 minutes of preparation should be prepared with cold infusion solutions. Limited Distribution product distributed by CVS Specialty. Prior to reconstitution, store at ≤25° C. Following reconstitution, may be stored at room temperature for 24 hours, although the manufacturer recommends use begin within 3 hours of reconstitution. Do Not refrigerate reconstituted solution. When stored at 30° C in polypropylene infusion pump syringes, deferoxamine 250 mg/mL in sterile water for injection retained 95% of initial concentration for 14 days	Yes	No
Deferoxamine (Desferal)	Iron Overload	Injection IM or IV	No	No	No	None	Yes	Yes
Dofetilide (Tikosyn)	Cardiac Disorders	Oral	Yes	No	No	None	No	Yes
Duopa	Movement Disorders	Enteral Suspension	No	No	Yes	Limited Distribution product distributed Not by CVS Specialty	Yes	No
Dupixent	Atopic Dermatitis	Injection SubQ	Yes	Yes	Yes	N/A	No	No
Dysport	Batten Disease	Injection IM	No	No	Yes	Refrigerate, protect from light	No	No
Egrifta	HIV Medications	Injection SubQ	Yes	Yes	No	Limited Distribution product distributed by CVS Specialty	Yes	No
						Limited Distribution product distributed by CVS Specialty. Refrigerate vials. Protect from light, do not freeze or shake. Should be used immediately after dilution in NS. However, solution for infusion may be stored under refrigeration for up to 24 hours.		
Elaprase	Lysosomal Storage Disorders	IV	No	No	No		Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Ellelyso Eligard (Canadian Brand name, See Lupron)	Lysosomal Storage Disorders Hormonal Therapies	IV Injection SubQ	No No	No No	No Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate intact vials; protect from light. Do not freeze. If not used immediately, the reconstituted solution may be refrigerated for ≤24 hours (protect from light) or at room temp for ≤4 hours (without protection from light). The solution diluted in NS for infusion may be refrigerated for ≤24 hours; protect from light. In total, the reconstituted and diluted products can be stored for ≤24 hours. Do not freeze.	Yes No	No No
Eloctate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Prior to reconstitution, store refrigerated at 2°C to 8°C; do not freeze. May also store at room temperature up to 6 mon; do not return to refrigerator. Store in original package to protect from light. Use within 3 hours of reconstitution; do not refrigerate after reconstitution	Yes	No
Elspar	Oncology- Injectable	Injection IM or IV	No	No	N/A	Limited Distribution product distributed by CVS Specialty	Yes	No
Emflaza	Duchenne Muscular Dystrophy	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Store at 20°C to 25°C; excursions permitted between 15°C to 30°C.	Yes	No
Empliciti	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C (36°F to 46°F). Protect from light (store in the original packaging until use); do not freeze or shake. Solutions diluted for infusion in NS or D5W may be stored at 2°C to 8°C (36°F to 46°F) for up to 24 hours (protected from light). A maximum of 8 hours of the 24 hour storage time may be at room temperature and room light. Infusion must be completed within 24 hours of lyophilized powder reconstitution.	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Enbrel	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate at 2°C to 8°C. Do Not shake. Do Not freeze or store in extreme heat or cold. Store in the original carton to protect from light or physical damage. Individual autoinjectors, prefilled syringes, dose trays (containing multi-use vials and diluent syringes), or prefilled pens may be stored between 20°C and 25°C (68°F and 77°F) for a maximum single period of 14 days (Enbrel) with protection from light and sources of heat and humidity. Once an autoinjector, prefilled syringe, dose tray, or prefilled pen has been stored at room temperature, it should Not be placed back into the refrigerator; discard after 14 days (Enbrel). Refrigerate unopened vials. Retain in original package to protect from light. After reconstitution and dilution, the infusion solution may be re-refrigerated for up to 4 hours). Don't freeze. Discard any unused portion	No	Yes
Entyvio	Inflammatory Bowel Disease	IV	No	No	Yes	Dispense in original container	No	No
Epclusa	Hepatitis C	Oral	Yes	No	Yes		No	Yes
Epogen	Anemia	Injection SubQ or IV	Yes	Yes	Yes	Vials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light. Limited Distribution product distributed by CVS Specialty	No	Yes
Epoprostenol sodium (Flolan)	Pulmonary Arterial Hypertension	IV	No	No	No	Limited Distribution product distributed by CVS Specialty. Store intact vials refrigerated at 2°C to 8°C; do not freeze. Preparations in infusion containers are stable for up to 12 hours refrigerated at 2°C to 8°C and up to 8 hours at room temperature of 20°C to 25°C	Yes	No
Erbitux	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Keep bottle closed tightly to protect from moisture.	Yes	Yes
Erivedge	Oncology- Oral/Topical	Oral	Yes	No	Yes		Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Erwinaze	Oncology- Injectable	Injection IM or IV	No	No	Yes	Refrigerate. Protect from light. Within 15 minutes of reconstitution, withdraw appropriate volume for dose into a polypropylene syringe. Do not freeze or refrigerate reconstituted solution; discard if not administered within 4 hours.	No	No
Esbriet	Pulmonary Disorders- Others	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store refrigerated or at room temperature; do not freeze. Protect from light. If refrigerated, remove from refrigeration at least 20 to 30 minutes before use.	Yes	No
Euflexxa	Osteoarthritis	Injection Intra-articular	No	No	No	Limited Distribution product distributed by CVS Specialty. Store intact vials at 25°C; excursions are permitted between 15°C to 30°C. Protect from light (store in original container). The reconstituted solution is stable for 1 hour at room temperature or for 24 hours at 5°C. Solutions diluted in NS for infusion are stable for 4 hours at room temperature (in addition to the 1 hour at room temperature following reconstitution).	No	Yes
Evomela Exjade	Oncology- Injectable Iron Overload	IV Oral	No Yes	No No	Yes Yes	Protect from moisture. Store at 2°C to 8°C. Do Not freeze. Protect from light and store in the original carton until ready for use. The diluted solution may be stored at 2°C to 8°C for up to 24 hrs; do Not freeze. If not used immediately following reconstitution, refrigerate solution and use within 3 hours; do not freeze or shake solution.	Yes Yes	No No
Exondys	Duchenne Muscular Dystrophy	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Protect from light. Store in original container.	No	Yes
Extavia	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Limited Distribution product distributed by CVS Specialty. Refrigerate. Protect from light. Store in original container.	No	Yes
Eylea	Retinal Disorders	Injection Intravitreal	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate intact vials. Reconstituted solutions and solutions diluted in NS are stable for 24 hours at 2°C and 8°C.	Yes	No
Fabrazyme	Lysosomal Storage Disorders	IV	No	No	No		Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Farydak	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store blister pack in original carton. Protect from light. Store intact vials at ≤25°C; store in the original package to protect from light. Do not freeze. Following reconstitution, do not refrigerate and administer within 3 hours.	Yes	No
Feiba NF	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store intact vials at ≤25°C; store in the original package to protect from light. Do not freeze. Following reconstitution, do not refrigerate and administer within 3 hours.	Yes	No
Feiba VH	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store intact vials at ≤25°C; store in the original package to protect from light. Do not freeze. Following reconstitution, do not refrigerate and administer within 3 hours. Limited Distribution product distributed Not by CVS Specialty. Store at 20°C to 25°C; excursions permitted to 15°C to 30°C. Store Oral solution in original carton to protect from light; use contents within 35 days after first opening bottle; discard any remaining solution after 35 days.	Yes	No
Ferriprox	Iron Overload	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Firazyr	Hereditary Angiodema	Injection SubQ	Yes	Yes	Yes	Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture.	Yes	No
Firmagon	Hormonal Therapies	Injection SubQ	No	No	Yes	Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.	No	Yes
Flebogamma	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Don't freeze, heat, or shake. Store at 2°C to 25°C (36°F to 77°F). Keep in original carton to protect from light.	No	Yes
Flebogamma DIF	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Refrigerate. Avoid excessive or vigorous agitation; do Not shake. Discard if intact vial is left at room temperature for >12 hours prior to use.	No	Yes
Follistim AQ	Infertility	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	No	Yes
Folotyn	Oncology- Injectable	IV	No	No	Yes	Refrigerate; do Not freeze (discard if freezing occurs). Protect from light. Discard pen 28 days after first injection, even if it still contains some unused solution. Do Not use if solution is cloudy, colored, or contains solid particles.	Yes	No
Forteo	Osteoporosis	Injection SubQ	Yes	Yes	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Fusilev Fuzeon	Oncology- Injectable HIV Medications	IV Injection SubQ	No Yes	No Yes	No No	Limited Distribution product distributed by CVS Specialty  Don't freeze, heat, or shake. Store at ≤25°C (≤77°F). May store diluted solution under refrigeration at 2°C to 8°C for up to 24 hours if originally prepared in a sterile laminar air flow environment.	Yes No	No Yes
GamaSTAN S/D Gammagard Liquid (Canadian Brand)	Immune Deficiencies & Related Disorders Immune Deficiencies & Related Disorders	Injection IM Injection SubQ or IV	No No	No No	No N/A	Don't freeze, heat, or shake. Store at ≤25°C . May store diluted solution under refrigeration at 2°C to 8°C for up to 24 hours if originally prepared in a sterile laminar air flow environment.	Yes Yes	No No
Gammagard S/D	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Don't freeze, heat, or shake. Store at 2°C to 8°C; may be stored at ≤25°C for up to 6 months. Limited Distribution product distributed by CVS Specialty. Do not freeze, heat, or shake. Store at 2°C to 25°C. Keep in original carton to protect from light. Do not freeze or use if solution has been frozen.	Yes	No
Gammaked	Immune Deficiencies & Related Disorders	Injection SubQ or IV	No	No	Yes	Don't freeze, heat, or shake. Store at 2°C to 25°C. Keep in original carton to protect from light. Do not freeze or use if solution has been frozen.	Yes	No
Gammaplex	Immune Deficiencies & Related Disorders	IV	No	No	No	Don't freeze, heat, or shake. Store at 2°C to 25°C. Keep in original carton to protect from light. Do not freeze or use if solution has been frozen.	Yes	No
Gamunex-C Ganirelix	Immune Deficiencies & Related Disorders Infertility	Injection SubQ or IV Injection SubQ	No Yes	No Yes	Yes No	Protect from light Refrigerated at 2°C to 8°C; do Not freeze. The carton of ancillary supplies should be stored at 25°C. After dispensing, store vials at 25°C; once dispensed, vials must be used within 90 days. Once reconstituted, store at <25°C; do Not shake or freeze; use within 3hr	Yes No	No Yes (generic)
Gattex	Gastrointestinal Disorders- Other	Injection SubQ	Yes	No	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Gazyva	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C ; do not freeze or shake. Protect from light. Solutions diluted in NS for infusion should be used immediately. If not used immediately, the diluted solutions may be stored up to 24 hours at 2°C to 8°C followed by 48 hours (including infusion time) at room temperature of ≤30°C.	Yes	No
Gel-One	Osteoarthritis	Injection Intra-articular	No	No	Yes	Store below 25°C (77°F); do not freeze	No	No
Gelsyn-3	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 25°C; do not freeze	No	No
Gemzar	Oncology- Injectable	IV	No	No	Yes	Don't freeze	No	Yes
GeNotropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
Genvisc 850	Osteoarthritis	Injection Intra-articular	No	No	No	Limited Distribution product distributed by CVS Specialty. Store below 30°C; do not freeze. Store in original package to protect from light.	Yes	No
Gilenya	Multiple Sclerosis	Oral	Yes	No	Yes	Protect from moisture. Dispense in original bottle; protect from high humidity and light.	No	Yes
Gilotrif	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty.	No	No
Glassia	Alpha-1 Antitrypsin Deficiency	IV	No	No	Yes	Refrigerate. If needed, may store at 15°C to 30°C for up to 1 month. Avoid exposure to high temperatures; protect from intense light. Do not freeze. Discard if syringe freezes	Yes	No
Glatiramer acetate (Copaxone)	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Store powder refrigerated or at room temperature. Protect from light; do not freeze. Following reconstitution, multidose vials may be stored under refrigeration or at room temperature for up to 28 days.	No	Yes
Gonal-F	Infertility	Injection SubQ	Yes	Yes	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Granix	Neutropenia	Injection SubQ	Yes	Yes	No	Refrigerate prefilled syringes. Protect from light. Do not shake. May be removed from for a single period of up to 5 days. If not used within 5 days, the product may be returned to refrigerator up to the expiration date. Dispose of syringes if stored at room temperature for more than 5 days. Exposure to -1°C to -5°C for up to 72 hours and temperatures as low as -15°C to -25°C for up to 24 hours do not adversely affect stability. Discard unused product. Refrigerate. Limited Distribution product distributed by CVS Specialty	No	Yes
H.P. Acthar Gel	Hormonal Therapies / Seizure Disorders	Injection SubQ or IM	Yes	Yes	Yes	Refrigerate. Limited Distribution product distributed by CVS Specialty. Do not freeze. Store in original carton. Undiluted solutions in a syringe and solutions diluted in normal saline for infusion are stable for up to 4 hours at room temperature or up to 24 hours refrigerated at 4°C	Yes	No
Halaven	Oncology- Injectable	IV	No	No	Yes	Dispense in original container	Yes	Yes
Harvoni	Hepatitis C	Oral	Yes	No	Yes		No	Yes
Helixate FS	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Prior to reconstitution, store refrigerated; do not freeze. Do not refrigerate after reconstitution. May also be stored at room temperature up to 12 mon; do not return to refrigerator. Protect from extreme exposure to light during storage. Use within 3 hrs of reconstitution. Refrigerate; do not freeze. Use within 3 hours of reconstitution. Do not refrigerate after reconstitution, precipitation may occur. May also be stored at room temperature not to exceed 30°C	Yes	Yes
Hemofil M	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Refrigerate. Don't freeze or shake. Use within 6 hours of entering vial. Do not freeze. Solutions reconstituted	Yes	Yes
Hepagam B	Immune Deficiencies & Related Disorders	Injection IM or IV	No	No	No		Yes	Yes
Herceptin	Oncology- Injectable	IV	No	No	Yes		Yes	No
Hetlioz	Sleep Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed Not by CVS Specialty. Protect from light and moisture.	Yes	No



Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Hizentra	Immune Deficiencies & Related Disorders	Injection SubQ	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at ≤25°C. Keep in original carton to protect from light. Do not freeze or use if solution has been frozen.	Yes	Yes
Humantrope	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Protect from light. Refrigerate	No	No
Humate-P	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Products are stable for three years. For single use; discard any unused contents in vial. Store intact vials at ≤25°C; do not freeze. Once reconstituted, do not refrigerate and use within 3 hours.	Yes	No
Humira	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Store at 2°C to 8°C (36°F to 46°F) in original container to protect from light; do Not freeze. Do Not use if frozen even if it has been thawed. Do Not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C (77°F) for up to 14 days; discard if Not used within 14 days.	No	Yes
Hyalgan	Osteoarthritis	Injection Intra-articular	No	No	No	Limited Distribution product distributed by CVS Specialty. Store at 2°C to 8°C (36°F to 46°F). Protect from light.	No	Yes
Hycamtin	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store below 25°C; do not freeze. Store Hymovis in their original packages	Yes	No
Hymovis	Osteoarthritis	Injection Intra-articular	No	No	No	Refrigerate. Don't freeze or shake. May be exposed to room temperature for a cumulative 7 days	No	No
HyperHEP B	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Store at 2°C to 8°C; do not freeze.	No	Yes
HyppeRHO S/D	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Don't freeze, heat, or shake. Store at 2°C to 8°C for up to 36 months; may store at ≤25°C for up to 3 months during the first 24 months from the date of manufacture (after 3 months at room temperature, discard); do not return vial to refrigerator after it has been stored at room temperature.	No	No
HyQvia	Immune Deficiencies & Related Disorders	Infusion SubQ	No	No	No	Limited Distribution product distributed by CVS Specialty	Yes	No
Ibrance	Oncology- Oral/Topical	Oral	Yes	No	Yes		Yes	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Iclusig	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty Store in refrigerator or at room temperature from 2°C to 25°C. Do not freeze; store in package to protect from light. Reconstituted solution should be used immediately or within 4 hrs of preparation.	Yes	No
Idelvion	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution product distributed by CVS Specialty. Store intact vial at 2°C to 8°C; do Not freeze. After reconstitution, vials may be stored at room temp for up to 1 hr or in a refrigerator at for up to 4 hours. Protect from light. Discard any unused portion.	Yes	No
Ilaris	Cryopyrin-Associated Periodic Syndrome	Injection SubQ	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 15°C to 25°C (59°F to 77°F)	Yes	Yes
Iluvien	Retinal Disorders	Implant	No	No	Yes	Protect from moisture	Yes	No
Imantinib (Gleevec)	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Keep in original container until dispensing.	No	Yes
Imbruvica	Oncology- Oral/Topical	Oral	Yes	No	Yes	refrigerated at 2°C to 8°C	Yes	No
Imlygic	Oncology- Injectable	Injection Intralesional	No	No	Yes	Protect from light.	Yes	No
Increlex (IGF-1 Deficiency)	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	No
Inflectra	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis	IV	No	No	Yes	Refrigerate intact vials at 2°C to 8°C. Manufacturer recommends that solutions diluted in NS for infusion should be used within 3 hours of preparation. However, a stability study of infliximab 0.4 mg/mL prepared in NS in polyvinyl chloride (PVC) bags found no loss of biological activity when stored refrigerated at 4°C for up to 14 days	No	Yes
Ingrezza	Movement Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed Not by CVS Specialty	Yes	No
Inlyta	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Intron A	Hepatitis C / Oncology- Injectable	Injection SubQ or IM	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials under refrigeration at 2°C to 8°C; do not freeze. After reconstitution of powder for injection, product should be used immediately, but may be stored under refrigeration for ≤24 hours.	Yes	Yes
Iressa	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Istodax	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Room temp	Yes	No
Ixempra	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Protect from light. Reconstituted solution (in the vial) is stable for up to 1 hour at room temperature; infusion solution diluted in appropriate solution for infusion is stable for 6 hours at room temperature if a pH range of 6 to 9 is maintained (infusion must be completed within 6 hours).	Yes	No
Ixinity	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store 250 unit strength at 2°C to 8°C; store 500 unit, 1,000 unit, 1,500 unit, 2,000 unit, and 3,000 unit strength at 2°C to 25°C. Do not freeze. Keep vial in the carton and protect from light. Infuse reconstituted solution immediately or within 3 hours when stored at room temperature. Do not refrigerate after reconstitution.	Yes	No
Jadenu	Iron Overload	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Jakafi	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Jevtana	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 25°C; excursions permitted between 15°C and 30°C. Do not refrigerate. Do not prepare or administer in PVC-containing infusion containers or polyurethane infusion sets. The initial reconstituted solution (at 10 mg/mL) is stable for 30 min in the vial and solutions for infusion are stable for up to 8 hours at room temperature (includes the 1 hour infusion) or 24 hrs refrigerated (includes the 1 hour infusion).	Yes	Yes
Juxtapid	Lipid Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C. Do not freeze or shake intact vials, reconstituted solution, or solutions diluted for infusion. Reconstituted vials do not contain preservative and should be used immediately, although may be stored for up to 24 hours at 2°C to 8°C. Solutions diluted for infusion in NS should be used immediately, although may be stored at 2°C to 8°C for up to 24 hours prior to use. This storage time is additional to the time allowed for the reconstituted vials.	Yes	No
Kadcyla	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate and protect from light	Yes	No
Kalbitor	Hereditary Angiodema	Injection SubQ	Yes	Yes	Yes	Limited Distribution product Not distributed by CVS Specialty. Store at 20°C to 25°C; excursions permitted to 15°C to 30°C; after mixing the granules, the product is stable for 1 hour.	Yes	No
Kalydeco	Cystic Fibrosis	Oral	Yes	No	Yes		Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Kanuma	Lysosomal Storage Disorders	IV	No	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Store intact vials under refrigeration in original carton to protect from light. Do not shake or freeze. If not used immediately, reconstituted product may be stored up to 24 hours in the refrigerator.	Yes	No
Kevzara Keytruda	Rheumatoid Arthritis Oncology- Injectable	Injection SubQ	Yes	Yes	Yes	Refrigerate, Protect from light. Do Not freeze or shake. After removal from the refrigerator, use within 14 days or discard.	No	Yes
		IV	No	No	Yes		Yes	No
Kineret	Cryopyrin-Associated Periodic Syndrome / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate. Do Not freeze. Do Not shake. Protect from light. Discard any unused portion.	No	Yes
Kisqali	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store in the original package	No	Yes
Kitabis Pak	Cystic Fibrosis	Nebulized solution	Yes	No	No	Store under refrigeration at 2°C to 8°C. May be stored in foil pouch (opened or unopened) at room temperature of 25°C for up to 28 days. Protect from light. The colorless to pale yellow solution may darken over time if Not stored under refrigeration; however, the color change does Not affect product quality. Do Not use if solution has been stored at room temperature for >28 days.	No	No
Koate-DVI	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	N/A		Yes	No
Kogenate FS	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Prior to reconstitution, store	Yes	Yes
Korlym	Cushing's Syndrome	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	No
Kovaltry	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Prior to reconstitution, store Diluted solution may be stored up to 4 hours at 2°C to 8°C. Diluted solution is also stable for 4 hours at room temperature of 20°C to 25°C; Refrigeration is preferred. The diluted solution should be protected from light, Not frozen, and used within 4 hours of dilution.	Yes	No
Krystexxa	Gout	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	No	No
Kuvan	Phenylketonuria	Oral	Yes	No	Yes	Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F)	Yes	No
Kyleena	Contraceptives	IUD	No	No	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Kynamro	Lipid Disorders	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty Limited Distribution product distributed by CVS Specialty. Refrigerate. Store in original carton until use to protect from light. Reconstituted drug (in the vial or in a syringe) and preparations diluted for infusion in D5W are stable for 4 hours at room temperature or for 24 hours refrigerated at 2°C to 8°C (36°F to 46°F). Store intact vials at 2°C to 8°C; do not freeze. Keep in original carton to protect from light. Do not shake. Solutions diluted for infusion may be stored for up to 24 hours refrigerated and for an additional 4 hours at room temperature (infusion must be completed within this time frame). If refrigerated, allow infusion solution to reach room temperature prior to administration.	Yes	No
Kyprolis	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Prior to dilution in NS or D5W, refrigerate intact vials. Do not freeze. Do not shake; protect from light. Following dilution, store at room temperature or refrigerate; use within 8 hours	Yes	No
Lartruvo	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Prior to dilution in NS or D5W, refrigerate intact vials. Do not freeze. Do not shake; protect from light. Following dilution, store at room temperature or refrigerate; use within 8 hours	No	No
Lemtrada	Multiple Sclerosis	IV	No	No	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	Yes
Lenvima	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Letairis	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes	Refrigerate. Don't freeze or shake	Yes	No
Leukine	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Room temp	No	Yes
Leuprolide acetate (Lupron)	Hormonal Therapies	Injection SubQ or IM	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	No	Yes
Levoleucovorin calcium (see Fusilev)	Oncology- Injectable	IV	No	No	N/A	Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). Protect from light. Limited Distribution product Not distributed by CVS Specialty	Yes	No
Liletta	Contraceptives	IUD	No	No	Yes	Specialty	Yes	No
Lonsurf	Oncology- Oral/Topical	Oral	Yes	No	Yes	If stored outside the original bottle, discard Orals after 30 days	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Lucentis	Retinal Disorders	Injection Intravitreal	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Keep prefilled syringe in sealed tray until ready to use.	Yes	No
Lumizyme	Lysosomal Storage Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials between 2°C and 8°C . Final solutions diluted in NS for infusion should be used immediately if possible but may be stored for up to 24; do not freeze.	Yes	No
Lupaneta Pack	Hormonal Therapies	Injection IM and Oral	No	No	Yes	Room temp	No	Yes
Lupron Depot	Hormonal Therapies	Injection IM	No	No	Yes	Room temp	No	Yes
Lynparza	Oncology- Oral/Topical	Oral	Yes	No	Yes	Orals: Store at 25°C; excursions permitted to 15°C to 30°C. Do not expose Orals to temperatures >40°C. Orals: Store at 20°C to 25°C; excursions permitted to 15°C to 30°C. Protect from moisture.	No	No
Macugen	Retinal Disorders	Injection Intravitreal	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 2°C to 8°C (36°F to 46°F); do Not freeze. Do Not shake vigorously.	Yes	No
Makena	Pre-Term Birth	Injection IM	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store refrigerated at 2°C to 8°C; do not freeze. Dispense in original bottle; do not remove desiccant. Protect from light and moisture. Do not transfer to pill boxes.	Yes	Yes
Mekinist	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate or room temp. Protect from light	Yes	Yes
MeNopur	Infertility	Injection SubQ	Yes	Yes	No	Store at 2°C to 8°C; do not freeze.	No	Yes
Micrhogam	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Limit Distribution products Not distributed by CVS Specialty	No	No
Mircera	Anemia	Injection SubQ or IV	Yes	Yes	No	Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). Limited Distribution product distributed by CVS Specialty	Yes	No
Mirena	Contraceptives	IUD	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Mitoxantrone (Novantrone)	Oncology- Injectable	IV	No	No	No	Limited Distribution product distributed by CVS Specialty	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
MoNoclate-P	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Refrigerate. Do not freeze. Use within 3 hrs of reconstitution. Do not refrigerate after reconstitution, precipitation may occur. May also be stored at 25°C for ≤6 mon. Store in original package to protect from light. When stored at refrigerator temperature, 2°C to 8°C, factor IX is stable for the period indicated by the expiration date on its label. Avoid freezing which may damage container for the diluent. May also be stored at room temperature not to exceed 25°C for up to 1 month. Reconstituted solution should be at room temperature and used within 3 hours of preparation.	Yes	No
MoNonine	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store below 25°C ]; do not freeze. Protect from light	Yes	No
MoNovisc	Osteoarthritis	Injection Intra-articular	No	No	Yes	Limited Distribution product distributed by CVS Specialty	No	Yes
Mozobil	Hematopoietics	Injection SubQ	No	No	No	Limited Distribution product distributed Not by CVS Specialty	No	No
Myalept	Lipodystrophy	Injection SubQ	Yes	Yes	Yes	Refrigerate, Protect from light, once diluted, use within 4 hr	Yes	No
Myobloc	Batten Disease	Injection IM	No	No	No	Refrigerate. Do not shake. Use within 6 hours of entering vial.	No	No
Nabi-HB	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Limited Distribution product distributed by CVS Specialty. Prior to reconstitution, refrigerate intact vials. Do not freeze. Do not shake. Protect from light. Following dilution in NS, use immediately. May store under refrigeration if used within 48 hours from the time of preparation to the completion of infusion. Do not store solution for infusion at room temperature. Allow vials to reach room temperature prior to dilution. Do not keep vials at room temperature >24 hours prior to dilution. Do not heat or microwave vials.	No	No
Naglazyme	Lysosomal Storage Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Natpara	Hormonal Therapies	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No



Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Neulasta	Neutropenia	Injection SubQ	Yes	Yes	Yes	Refrigerate. Don't freeze or shake. Protect from light	No	Yes
Neupogen	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate. Store in the original carton. Protect from light. Protect from direct sunlight. Avoid freezing, manufacturer recommends using within 24 hours due to the potential for bacterial contamination. Indicates Limited Distribution products distributed by CVS Specialty.	No	Yes
Nexavar	Oncology	Oral	Yes	No	Yes	Hazardous agent (NIOSH 2016 [group 2]).	Yes	No
NexplaNon	Contraceptives	Subdermal	No	No	Yes	Use appropriate precautions for receiving, handling, administration, and disposal. Gloves (single) should be worn during receiving, unpacking, and placing in storage. Limited Distribution product distributed by CVS Specialty	Yes	No
Ninlaro	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Do not freeze. Store in original packaging until immediately prior to use.	Yes	No
Noritropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate, but do Not freeze. Protect from light	No	No
Northera	Cardiac Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes
Novoeight	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Don't freeze. May also be stored at room temperature up to 12 months; do not return to refrigerator; after 12 months at room temp, use immediately or discard. Store in original package to protect from light. Use within 4 hours of reconstitution.	Yes	No
Novoseven RT	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Prior to reconstitution, store between 2°C to 25°C; do not freeze. Protect from light. Reconstituted solutions may be stored at room temperature or under refrigeration, but must be infused within 3 hours of reconstitution. Do not freeze reconstituted solutions. Do not store reconstituted solutions in syringes.	Yes	Yes
Nplate	Immune (Idiopathic) Thrombocytopenicpurpua	Injection SubQ	No	No	Yes	Refrigerate. Protect from light. Store in original carton until use.	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Nucala	Allergic Asthma	Injection SubQ	No	No	Yes	Limited Distribution producted distributed by CVS Specialty	Yes	Yes
Nuplazid	Movement Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Nutropin AQ	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Store unopened vials in the refrigerator, away from heat and direct light. Do not freeze. Should be used within 28 days.	No	No
Nuwiq	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Refrigerate. Don't freeze. May also be stored at room temp up to 3 months; do not return to refrigerator. Store in original package to protect from light. Use within 3 hrs of reconstitution	Yes	No
Obizur	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution producted distributed by CVS Specialty. Store refrigerated at 2°C to 8°C. Do not freeze. Store the original package to protect from light. Use within 3 hours after reconstitution; discard any unused solution if not used within 3 hours after reconstitution.	Yes	No
Ocaliva	Gastrointestinal Disorders- Other	Oral	Yes	No	Yes	-	Yes	No
Ocrevus	Multiple Sclerosis	IV	No	No	Yes	Refrigerate in the outer carton to protect from light. Do not freeze. Do not shake. Solutions diluted for infusion should be used immediately, however may be stored for up to 24 hours refrigerated and 8 hours at room temperature (including infusion time); discard if infusion cannot be completed on the same day.	No	No
Octagam	Immune Deficiencies & Related Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 2°C to 25°C (36°F to 77°F).	Yes	Yes
Octreotide acetate (Sandostatin)	Acromegaly	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate	No	Yes
Odomzo	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	No
Ofev	Pulmonary Disorders- Others	Oral	Yes	No	Yes	Limited Distribution producted distributed Not by CVS Specialty Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Protect from high humidity and avoid excessive heat.	Yes	No
Olysio	Hepatitis C	Oral	Yes	No	Yes	Store in original bottle and protect from light	No	Yes
Omnitrope	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Protect from light. Refrigerate	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Oncaspar	Oncology- Injectable	Injection IM or IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Do not shake; protect from light.	Yes	No
Onivyde	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Refrigerate. Do not freeze. Protect from light.	Yes	No
Opdivo	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Protect from light. Do not shake.	Yes	No
Opsumit	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Oralair	Allergen ImmuNotherapy	Oral	Yes	No	No	Limited Distribution products distributed by CVS Specialty. Store between 20°C and 25°C (68°F and 77°F); excursions are permitted between 15°C and 30°C (59°F and 86°F). Protect from moisture.	Yes	No
Orenica	Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Refrigerate, protect from light, must used within 24 hr of reconstituting	No	No
Orenitram	Pulmonary Arterial Hypertension	Oral	Yes	No	N/A	Limited Distribution product distributed by CVS Specialty	Yes	No
Orfadin	Lysosomal Storage Disorders	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Refrigerate	Yes	No
Orkambi	Cystic Fibrosis	Oral	Yes	No	No	Limited Distribution product distributed by CVS Specialty	Yes	No
Orthovisc	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 25°C ]; do not freeze. Protect from light	No	Yes
Otezla	Psoriasis	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes
Otrexup	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	No	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Refrigerate or room temp for 30 days. Protect from light	No	Yes
Ovidrel	Infertility	Injection SubQ	Yes	Yes	No		No	Yes
Oxaliplatin	Oncology- Injectable	IV	No	No	No	Refrigerate or room temp for 30 days. Protect from light	No	Yes
Ozurdex	Retinal Disorders	Implant	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 15°C to 30°C (59°F to 86°F).	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Pegasys	Hepatitis C	Injection SubQ	Yes	No	Yes	Refrigerate and protecting from light Intact vial: May be stored at room temperature for up to 14 days  Prefilled syringe: May be stored at room temperature for up to 6 days  Refrigerate. Protect from light. Do not freeze. Do not shake. Solutions diluted for infusion in NS should be used immediately; if not used immediately, may be stored at 2°C to 8°C for up to 24 hours.	No	Yes
Perjeta	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store in the closed original carton to protect from light, between 2°C and 8°C. Do not freeze; discard if frozen.  If refrigeration is unavailable, may store between 2°C and 25°C for up to 30 days, protected from light. May remove from, and return to, a refrigerator if necessary. The total combined time out of refrigeration, within a temperature range of 2°C to 25°C, should not exceed 30 days.	Yes	No
Plegridy	Multiple Sclerosis	Injection SubQ	Yes	Yes	No	Limited Distribution product distributed by CVS Specialty	Yes	No
Pomalyst	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty.  Refrigerate. Do not freeze. Store in original carton to protect from light. Do not shake. Solutions diluted for infusion in NS are stable for up to 4 hours at room temperature or 24 hours at 2°C to 8°C; do not freeze or shake.	Yes	No
Portrazza	Oncology- Injectable	IV	No	No	Yes	Refrigerate in the outer carton to protect from light. Time out of refrigeration should Not exceed 24 hours at 25°C. Do Not freeze. Do Not expose to extreme heat.	Yes	No
Praluent	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Do Not shake.  Prior to use, refrigerate. Once diluted, may be stored at 2°C to 8°C for 24 hours; refrigerate during transit. Do Not freeze. Protect from light.	No	Yes
Prialt	Pain Management	Injection Intrathecal	No	No	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Privigen	Immune Deficiencies & Related Disorders	IV	No	No	No	Limited Distribution product distributed by CVS Specialty. Do not freeze, heat, or shake. Store at ≤25°C. Do not freeze or use if solution has been frozen. Protect from light.	Yes	Yes
Procrit	Anemia	Injection SubQ or IV	Yes	Yes	No	Viials should be stored at 2°C to 8°C (36°F to 46°F). Do Not freeze. Do Not shake. Protect from light. Limited Distribution product Not distributed by CVS Specialty. Protect from light and moisture. Dispense delayed-release Orals in original package; do not subdivide or repackage.	No	Yes
Procysbi	Lysosomal Storage Disorders	Oral	Yes	No	Yes	Storage temperature should not exceed 25°C; do not freeze.	Yes	No
Profilnine SD	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limit Distribution products Not distributed by CVS Specialty	Yes	No
Prolastin-C	Alpha-1 Antitrypsin Deficiency	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials under refrigeration at 2°C to 8°C. Protect from light. Plastic (polyvinyl chloride) bags result in more consistent drug delivery and are recommended. According to the manufacturer, reconstituted vials and solutions diluted in D5W for infusion are stable for 48 hours at room temperature or refrigerated although refrigeration is preferred because they do not contain preservatives. Do not freeze.	Yes	No
Proleukin	Oncology- Injectable	IV	No	No	Yes	Refrigerate in original carton. Do Not freeze. Prior to use, bring to room temperature of 25°C in original container (usually takes 15 to 30 minutes); do Not use any other methods for warming. Use within 14 days once at room temperature. Protect from direct heat and light; do Not expose to temperatures >25°C.	Yes	Yes
Prolia	Osteoporosis Immune (Idiopathic)	Injection SubQ	No	No	Yes	Avoid vigorous shaking. Room temp. Dispense in original bottle.	No	Yes
Promacta	Thrombocytopenicpurpua	Oral	Yes	No	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Provenge	Oncology- Injectable	IV	No	No	Yes	Do not remove the infusion bag from the insulated polyurethane container within the shipping box until administration (do not remove the insulated container from the shipping box, or open the lid of the insulated container, until administration). Product may only remain at room temperature for ≤3 hours once removed from shipping container; after removal from shipping container, do not return product to container. Infusion must begin prior to product expiration. Store at 2°C to 8°C in protective foil to protect from light. Refrigerate during transport and do not expose to room temp	No	No
Pulmozyme	Cystic Fibrosis	Nebulized Solution	Yes	No	Yes	for ≥24 hours. Limited Distribution product distributed by CVS Specialty. Orals: Store at 15°C to 25°C. Store in a dry place. Suspension: Store at 15°C to 25°C. Do not store above 25°C. Store in a dry place. Use within 8 weeks after opening.	No	Yes
Purixan	Oncology- Oral/Topical	Oral	Yes	No	N/A	Limit Distribution products Not distributed by CVS Specialty. Store at ≤25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F). Protect from light. Once overwrap package is opened, use within 24 hours.	Yes	No
Radicava	Amyotrophic Lateral Sclerosis	IV	No	No	Yes	Store intact vials and autoinjectors between 20°C and 25°C (68°F and 77°F); excursions may be permitted between 15°C and 30°C (59°F and 86°F). Protect from light. Limited Distribution product distributed by CVS Specialty.	Yes	No
Rasuvo	Psoriasis / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate. Refrigerate; do not freeze. Protect from heat and light. Refrigeration is preferred; however, if needed, may be stored at 2°C to 25°C for up to 30 days.	No	Yes
Ravicti Rebetol Solution	Urea Cycle Disorders Hepatitis C	Oral Oral	Yes Yes	No No	Yes No		Yes No	No No
Rebif	Multiple Sclerosis	Injection SubQ	Yes	Yes	Yes		No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Recombinate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	May also be stored at room temperature, not to exceed 30°C (86°F). Use within 3 hours of reconstitution.	Yes	Yes
Remicade	Inflammatory Bowel Disease / Psoriasis / Rheumatoid Arthritis	IV	No	No	Yes	Refrigerate intact vials at 2°C to 8°C. Manufacturer recommends that solutions diluted in NS for infusion should be used within 3 hours of preparation. However, a stability study of infliximab 0.4 mg/mL prepared in NS in polyvinyl chloride (PVC) bags found no loss of biological activity when stored refrigerated at 4°C for up to 14 days	No	Yes
Remodulin	Pulmonary Arterial Hypertension	Injection SubQ or IV	No	No	No	Limited Distribution product distributed by CVS Specialty Refrigerate in the original carton. May also be kept at room temperature in the original carton; however, must use within 30 days (discard if Not used within 30 days). Protect from direct light and do Not expose to temp above 25°C. Do Not freeze. Do Not shake.	Yes	No
Repatha	Lipid Disorders- PSK9 Inhibitors	Injection SubQ	Yes	Yes	Yes	Refrigerate or room temp. Protect from light	No	Yes
Repronex	Infertility	Injection SubQ or IM	Yes	Yes	No	Limited Distribution product distributed by CVS Specialty.	No	Yes
Retisert	Retinal Disorders	Implant	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Revlimid	Oncology- Oral/Topical Immune Deficiencies & Related Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes
Rhogam	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Store at 2°C to 8°C (35°F to 46°F); do not freeze. Store at 2°C to 8°C; do not freeze. Protect from light.	No	Yes
Rhophylac	Immune Deficiencies & Related Disorders	Injection IM or IV	No	No	No	Store at 2°C to 25°C in original carton; do not freeze. Protect from light. RiaSTAP is stable for 8 hours after reconstitution when stored at 20°C to 25°C ; Discard partially used vials.	No	Yes
Riastap	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store at 25°C (77°F); excursions permitted between 15°C and 30°C (59°F and 86°F)	Yes	No
Ribapak	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Ribasphere	Hepatitis C	Oral	Yes	No	No	None	No	Yes
Ribavirin caps (rebetol)	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes
Ribavirin tabs (Copegus, Moderiba)	Hepatitis C	Oral	Yes	No	Yes	None	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Rituxan	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Do not shake. Protect vials from direct sunlight. Solutions for infusion in NS or D5W are stable at 2°C to 8°C for 24 hours and at room temperature for an additional 24 hours (although because there is no preservative, the manufacturer recommends storing refrigerated). Store at 2°C to 8°C; do not freeze. May store at room temperature not to exceed 30°C for up to 36 months; do not return to refrigerator. Infuse at room temperature and within 3 hours of reconstitution; do not refrigerate after reconstitution	Yes	No
Rixubis	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	none	Yes	No
Rubraca	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	No	No
Ruconest	Hereditary Angiodema	IV	Yes	Yes	Yes	Store in the original package to protect from moisture.	Yes	No
Rydapt	Oncology- Oral/Topical	Oral	Yes	No	Yes	Hazardous Agent	No	Yes
Sabril	Seizure Disorders	Oral	Yes	No	Yes	Protect from light.	Yes	No
Saizen	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Refrigerate	No	Yes
Samsca	Electrolyte Disorders	Oral	Yes	No	Yes	Store at 25°C; excursions permitted between 15°C and 30°C.	No	No
Sandostatin Lar	Acromegaly	Injection IM	No	No	Yes	Refrigerate	No	Yes
Sensipar	Renal Disease	Oral	Yes	No	Yes	Store at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F).	No	Yes
Serostim	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	No	Limited Distribution product distributed by CVS Specialty. Can store in refrigeration after reconstitution	Yes	No
Signifor	Cushing's Syndrome	Injection SubQ	Yes	Yes	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	No
Signifor Lar	Acromegaly	Injection IM	No	No	Yes	Limit Distribution products Not distributed by CVS Specialty	Yes	No
Sildenafil (Revatio)	Pulmonary Arterial Hypertension	Oral	Yes	No	No		No	Yes
Simponi	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ	Yes	Yes	Yes	Refrigerate. Do not freeze. Do not shake. Store product in original carton to protect from light.	No	Yes



Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Simponi Aria	Rheumatoid Arthritis	IV	No	No	Yes	Refrigerate; do Not freeze. Do Not shake. Store product in original carton to protect from light IV: Solutions diluted for infusion may be stored at room temperature for 4 hours.	No	Yes
Skyla	Contraceptives	IUD	No	No	Yes	Store at 25°C (77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). Limited Distribution product distributed by CVS Specialty	Yes	No
Sodium phenylbutyrate (Buphenyl)	Urea Cycle Disorders	Oral	Yes	No	Yes	Store at 15°C to 30°C (59°F to 86°F). Limited Distribution product distributed by CVS Specialty	Yes	No
Soliris	Paroxysmal Nocturnal Hemoglobinuria	IV	No	No	Yes	Refrigerate	Yes	No
Somatuline Depot	Acromegaly	Injection SubQ	No	No	Yes	Limited Distribution products distributed by CVS Specialty	No	No
Somavert	Acromegaly	Injection SubQ	Yes	Yes	Yes	Dispense in original container	Yes	No
Sovaldi	Hepatitis C	Oral	Yes	No	Yes	Protect from light. Refrigerate. Discard after 30 hours after being out of original container	No	Yes
Spinraza	Spinal Muscular atrophy	Injection Intrathecal	No	No	No	none	No	No
Sprycel	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate in original container to protect from light; do not freeze. Do not use if frozen even if it has been thawed. Do not store in extreme heat or cold. If needed, may be stored at room temperature up to a maximum of 25°C for up to 14 days; discard if not used within 14 days.	No	Yes
Stelara	Inflammatory Bowel Disease / Rheumatoid Arthritis	Injection SubQ or IV	Yes	Yes	Yes	Store at room temperature not to exceed 25°C. Discard 6 months after opening bottle. Store bottle in upright position.	No	Yes
Stimate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	Nasal Solution	No	No	No	Limited Distribution product distributed by CVS Specialty. Store Orals in the original bottle and protect from moisture (do not remove the desiccant); keep container tightly closed. Discard any unused Orals 7 weeks after opening the bottle.	Yes	No
Stivarga	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes
Strensiq	Electrolyte Disorders	Injection SubQ	Yes	No	Yes	Not distributed by CVS Specialty	Yes	No
Supartz FX	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 30°C (86°C); c	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Supprelin LA Sudent	Hormonal Therapies Oncology- Oral/Topical	Implant Oral	No Yes	No No	Yes Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Keep implant wrapped in the amber pouch for protection from light; do Not freeze. Do Not open implant vial until just before the time of insertion. The implantation insertion kit should be stored at room temperature (do Not refrigerate insertion kit). none	Yes No	No Yes
Sylatron	Oncology- Injectable	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty. Prior to reconstitution, store Redipen at 2°C to 8°C. Store intact vials at 25°C; excursions permitted to 15°C to 30°C. Do not freeze. Once reconstituted each product should be used immediately or may be stored for ≤24 hours at 2°C to 8°C; do not freeze. Do not shake. Keep away from heat. Products do not contain preservative (single use; do not freeze).	Yes	Yes
Sylvant	Oncology- Injectable	IV	No	No	Yes	Store intact vials at 2°C to 8°C; protect from light. Reconstituted solution should be further diluted in D5W for infusion within 2 hours; complete infusion within 4 hours of dilution of the reconstituted solution to the infusion container. Discard any unused portion of the reconstituted solution or solution diluted for infusion.	No	No
Synagis	Respiratory Syncytial Virus	Injection IM	No	No	Yes	Refrigerate in original container. Don't freeze Limited Distribution product Not distributed by CVS Specialty. Protect from light (intact vial and reconstituted solutions). Reconstituted solution should be used within 12 hours if stored at room temperature or within 6 days (144 hours) if refrigerated	No	No
Synribo	Oncology- Injectable	Injection SubQ	No	No	Yes	refrigerated	Yes	No
Synvisc	Osteoarthritis	Injection Intra-articular	No	No	No	Store below 30°C (86°C); c	No	Yes
Synvisc One	Osteoarthritis	Injection Intra-articular	No	No	Yes	Store below 30°C (86°C); c	No	Yes
Tafinlar	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Tagrisso	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Taltz	Psoriasis	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty Refrigerate Limited Distribution product distributed by CVS Specialty Store at 25°C (77°F); excursions are permitted between 15°C to 30°C (59°F to 86°F). An Oral suspension (10 mg/mL) prepared using erlotinib 150 mg Orals and Ora-Plus:Ora-Sweet (1:1 vehicle) is stable for at least 28 days at room temperature; do not refrigerate due to potential increased viscosity	Yes	Yes
Tarceva	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	Yes	No
Targretin GelY	Oncology- Oral/Topical	Gel	Yes	No	Yes		No	Yes
Tasigna	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store intact vials between 2°C to 25°C. Protect from bright light. Freezing does not adversely affect the product. Solutions diluted for infusion in D5W or NS in non-PVC containers should be used within 6 hours of preparation, including infusion time, when stored between 2°C to 25°C or within 48 hours when stored between 2°C to 8°C	No	Yes
Taxotere	Oncology- Injectable	IV	No	No	No	Refrigerate. Do not freeze. Do not shake. Store in original carton to protect from light. Solutions diluted in NS for infusion should be used immediately after preparation; if not used immediately, may be stored for up to 6 hours (including administration time) at room temperature or 24 hours refrigerated at 2°C to 8°C. Do not freeze.	No	No
Tecentriq	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Protect Orals from light and store in the original container. Dispense in original container	No	No
Tecfidera	Multiple Sclerosis	Oral	Yes	No	Yes		Yes	No
Technivie	Hepatits C	Oral	Yes	No	Yes		No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Temodar Temozolomide (Temodar)	Oncology- Injectable Oncology- Oral/Topical	Oral or IV Oral	No	No	Yes	Store intact vials refrigerated at 2°C to 8°C. Reconstituted vials may be stored for up to 14 hours at room temperature of 25°C; infusion must be completed within 14 hours of reconstitution.	No	Yes
			Yes	No	Yes		none	Yes
Tetrabenazine (Xenazine)	Movement Disorders	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Prior to reconstitution, store at 2°C to 8°C (36°F to 46°F). Following reconstitution with the provided diluent, store under refrigeration and use within 14 days (5 mg vial) or 28 days (10 mg vial); do Not freeze.	Yes	No
Tev-Tropin	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	No	Protect from light. Keep in original package.	No	No
Thalomid	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C. Protect from light. May store reconstituted solution for up to 24 hours between 2°C and 8°C; avoid microbial contamination. If reconstituted solution is not refrigerated, use within 3 hours. Discard unused portion of the vial.	No	Yes
Thyrogen	Oncology- Injectable	Injection IM	No	No	Yes	Limited Distribution product distributed by CVS Specialty.	Yes	No
Tobi Podhaler	Cystic Fibrosis	Nebulized Solution	Yes	No	No	Store in original package at 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F). Protect from moisture.	Yes	Yes
Tobramycin inhalation solution (Tobi)	Cystic Fibrosis	Nebulized Solution	Yes	No	No	Limited Distribution product distributed by CVS Specialty. Store intact vials refrigerated at 2°C to 8°C. Diluted solution in the vial (10 mg/mL) is stable for 24 hours at room temperature. Solutions diluted for infusion (in NS) must be infused within 6 hours of preparation. Protect from light during storage, preparation, and handling.	No	No
Torisel	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty.	Yes	Yes
Tracleer	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty.	Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Treanda	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Solution: Store intact vials between 2°C to 8°C; protect from light. Solutions diluted for infusion in NS or D2.51/2NS are stable for up to 24 hours when stored at 2°C to 8°C or for up to 2 hours when stored at 15°C to 30°C and room light. Infusion must be completed within these time frames. Store at 20°C to 25°C. Do Not freeze MIXJECT system. Administer immediately after reconstitution.	Yes	Yes
Trelstar	Hormonal Therapies	Injection IM	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Prior to reconstitution, store refrigerated at 2°C to 8°C. Do not freeze; protect from light. If the reconstituted solution is not used immediately, store refrigerated or at room temperature not to exceed 25°C for up to 3 hours; discard after 3 hours.	No	No
Tretten	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Limited Distribution product distributed by CVS Specialty. Store refrigerated. Do Not freeze or heat. After first use, store for up to 30 days at 20°C to 25°C	Yes	No
Tykerb	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store concentrated solution under refrigeration; do not freeze. Protect from light. Do not shake. Following dilution, may store refrigerated for use within 8 hours.	Yes	Yes
Tymlos	Osteoporosis	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C; do not freeze. Do not shake. Keep the vial in the outer carton to protect from light. Solutions diluted for infusion in NS should be stored at 2°C to 8°C. Initiate infusion within 4 hours of preparation. Discard diluted solution 24 hours after preparation.	No	No
Tysabri	Inflammatory Bowel Disease / Multiple Sclerosis	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C; do not freeze. Do not shake. Keep the vial in the outer carton to protect from light. Solutions diluted for infusion in NS should be stored at 2°C to 8°C. Initiate infusion within 4 hours of preparation. Discard diluted solution 24 hours after preparation.	Yes	No
Tyvaso	Pulmonary Arterial Hypertension	Nebulized Solution	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Store intact vials at 2°C to 8°C; do not freeze. Do not shake. Keep the vial in the outer carton to protect from light. Solutions diluted for infusion in NS should be stored at 2°C to 8°C. Initiate infusion within 4 hours of preparation. Discard diluted solution 24 hours after preparation.	Yes	No
Unituxin	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C; do not freeze. Do not shake. Keep the vial in the outer carton to protect from light. Solutions diluted for infusion in NS should be stored at 2°C to 8°C. Initiate infusion within 4 hours of preparation. Discard diluted solution 24 hours after preparation.	Yes	No
Upravi	Pulmonary Arterial Hypertension	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store intact vials at 2°C to 8°C; do not freeze. Do not shake. Keep the vial in the outer carton to protect from light. Solutions diluted for infusion in NS should be stored at 2°C to 8°C. Initiate infusion within 4 hours of preparation. Discard diluted solution 24 hours after preparation.	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Valchlor	Oncology- Oral/Topical	Gel	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Prior to dispensing, store in freezer at -25°C to -15°C. After dispensing, refrigerate at 2°C to 8°C; apply immediately (or within 30 minutes) after removal from refrigerator; return to refrigerator promptly after each use. Discard unused product 60 days after opening.	Yes	No
Valstar	Oncology- Injectable	Injection Intravesically	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Do not freeze. Solutions diluted in 0.9% sodium chloride are stable for 12 hours at room temperature. Refrigerate. Keep implant wrapped in the amber pouch for protection from light; do Not freeze. Do Not open implant vial until just before the time of insertion. The implantation insertion kit should be stored at room temperature (do Not refrigerate insertion kit).	Yes	No
Vantas	Hormonal Therapies	Implant	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Store at 2°C to 8°C; do not freeze. Discard any unused portion.	No	No
Varizig	Immune Deficiencies & Related Disorders	Injection IM	No	No	No	Limited Distribution product distributed by CVS Specialty. Store intact vials in the original carton at 2°C to 8°C until the time of use. Do not freeze; do not shake. Protect from direct sunlight. Solutions diluted in NS for infusion should be used within 6 hours of preparation if stored at room temperature or within 24 hours of dilution if stored at 2°C to 8°C; do not freeze.	Yes	No
Vectibix	Oncology- Injectable	IV	No	No	Yes		Yes	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Velcade	Oncology- Injectable	Injection SubQ or IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Prior to reconstitution, store intact vials at 25°C; excursions are permitted between 15°C and 30°C. Once reconstituted, the manufacturer recommends use within 8 hours of reconstitution. However, stability studies have demonstrated solutions of 1 mg/mL (vial or syringe) may be stored at room temperature for up to 3 days, or under refrigeration for up to 5 days; or refrigerated in the original vial for up to 15 days. Protect from light. After preparation, keep bortezomib in a location away from the separate storage location recommended for intrathecal medications.	Yes	Yes
Velettri	Pulmonary Arterial Hypertension	IV	No	No	No	Limited Distribution product distributed by CVS Specialty. Refrigerated	Yes	No
Venclexta	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Store at or below 30°C (86°F).	Yes	No
Ventavis	Pulmonary Arterial Hypertension	Nebulized Solution	Yes	No	No	Limited Distribution product distributed by CVS Specialty	Yes	No
Viekira Pak	Hepatits C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Viekira XR	Hepatits C	Oral	Yes	No	Yes	Dispense in original container	No	Yes
Vimizim	Lysosomal Storage Disorders	IV	No	No	N/A	Limited Distribution product distributed by CVS Specialty	Yes	No
Vistogard	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	No
Visudyne	Retinal Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Injection: Store uNopened kit at 2°C to 8°C (36°F to 46°F). Kit may be kept at room temperature of ≤25°C (77°F) for ≤7 days prior to use; do Not freeze. Following reconstitution of the suspension, administer immediately.	Yes	No
Vivitrol	Alcohol/Opioid Withdrawal	Injection IM	No	No	Yes	Oral: Store at 20°C to 25°C (68°F to 77°F).	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Vonvendi	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store intact vials at 2°C to 8°C in original box; do not freeze. Protect from extreme exposure to light. Intact vials may also be stored at room temperature up to 30°C (86°F) for ≤12 months. Once stored at room temperature, do not return to refrigerator. Use reconstituted solution immediately; or may store at room temperature ≤27°C for up to 3 hours. Do not refrigerate after reconstitution.	Yes	No
Votrient	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	Yes
Vpriv	Lysosomal Storage Disorders	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate intact vials. Once reconstituted, the product should be used immediately. If immediate use is not possible, the reconstituted or diluted product may be stored for up to 24 hours at 2°C to 8°C. The infusion should be completed within 24 hours of reconstitution. Do not freeze. Protect from light. Discard any unused solution.	Yes	No
Wilate	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Store intact vials under refrigeration at 2°C to 8°C; do not freeze. Store in original container to protect from light. Intact vials may also be stored at room temperature for ≤6 months. Once stored at room temperature, do not return to the refrigerator. Following reconstitution, use solution immediately. Store at 2°C to 8°C ]; do not freeze.	Yes	No
Wunrho SDF	Immune Deficiencies & Related Disorders	Injection IM or IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	No	Yes
Xalkori	Oncology- Oral/Topical	Oral	Yes	No	Yes	Store at room temp	Yes	No
Xeljanz	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeljanz XR	Rheumatoid Arthritis	Oral	Yes	No	Yes	Store at room temp	No	Yes
Xeomin	Batten Disease	Injection IM	No	No	Yes	After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) and administer within 24 hours. Vial should only be used for one injection session and one patient.	No	No
Xermelo	Gastrointestinal Disorders- Other	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	No



Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Xgeva	Oncology- Injectable	Injection SubQ	No	No	Yes	Refrigerate. Do not freeze. Prior to use, bring to room temperature of 25°C in original container (usually takes 15 to 30 minutes); do not use any other methods for warming. Use within 14 days once at room temperature. Protect from direct heat and light; do not expose to temperatures >25°C. Avoid vigorous shaking. Limited Distribution product Not distributed by CVS Specialty. Prior to use, store lyophilized powder and diluent refrigerated; do Not freeze. Once reconstituted, may be stored at room temperature for ≤1 hour or refrigerated for ≤4 hrs	No	Yes
Xiaflex	Dupuytren's Contracture	Injection Intralesional	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Xolair	Allergic Asthma	Injection SubQ	No	No	Yes	CVS Specialty Limited Distribution product distributed by CVS Specialty. Protect from moisture; keep bottle tightly closed.	Yes	No
Xtandi	Oncology- Oral/Topical	Oral	Yes	No	Yes	Refrigerate. Don't freeze. May also be stored at room temperature up to 3 months; after room temperature storage, product may be returned to the refrigerator until the expiration date; however, do not store at room temperature and return to refrigerator temperature more than once. Avoid prolonged exposure to light during storage. Use within 3 hours of reconstitution. Limited Distribution product distributed by CVS Specialty.	Yes	No
Xyntha	Hemophilia, Von Willebrand Disease, & Related Bleeding Disorders	IV	No	No	No	Refrigerate. do not freeze. Protect from light. Prior to preparation, allow vials to sit at room temperature for ~5 minutes. Solutions diluted for infusion in NS or D5W are stable for up to 24 hours refrigerated or at room temperature.	Yes	No
Yervoy	Oncology- Injectable	IV	No	No	Yes		Yes	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Yondelis	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Solutions diluted for infusion in NS or D5W should be used within 30 hours of reconstitution (infusion should be completed within that 30 hours).	Yes	No
Zaltrap	Oncology- Injectable	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Protect from light (store in original outer carton). Solutions diluted for infusion in D5W or NS may be stored in refrigerator for up to 24 hours, or at 20°C to 25°C for up to 8 hours.	Yes	No
Zarxio	Neutropenia	Injection SubQ or IV	Yes	Yes	No	Refrigerate in the original carton. Protect from light. Avoid freezing; if frozen, thaw in the refrigerator before administration. Discard if frozen more than once. Do not shake. Transport via a pneumatic tube has not been studied. Solutions diluted for infusion may be stored at room temperature for up to 24 hours (infusion must be completed within 24 hours of preparation).	No	No
Zavesca	Lysosomal Storage Disorders	Oral	Yes	No	Yes	Note: Extended storage information may be available; contact product manufacturer to obtain current recommendations.	Yes	No
Zejula	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty	Yes	No
Zelboraf	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product distributed by CVS Specialty. Store in the original container with the lid tightly closed.	Yes	No
Zemaira	Alpha-1 Antitrypsin Deficiency	IV	No	No	Yes	Limited Distribution product distributed by CVS Specialty	Yes	No
Zepatier	Hepatits C	Oral	Yes	No	Yes	Keep in original blister pack until time of use; protect from moisture.	No	Yes

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Zinbryta	Multiple Sclerosis	Injection SubQ	Yes	Yes	Yes	Limited Distribution product distributed by CVS Specialty. Refrigerate. Do not expose to temperatures above 30°C. Do not freeze (discard if frozen). Keep in the original carton to protect from light. If refrigeration is unavailable, may be stored protected from light for up to 30 days at up to 30°C. Do not place back into the refrigerator after allowing it to warm to room temperature. Discard after 30 days without refrigeration. Store at room temp. Keep in foil pouch until ready to use to protect from light and moisture.	Yes	No
Zoladex	Hormonal Therapies	Implant	No	No	Yes	Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration. Store concentrate vials and ready-to-use bottles at 25°C; excursions permitted to 15°C to 30°C. Diluted solutions for infusion in D5W or NS which are not used immediately after preparation should be refrigerated at 2°C to 8°C. Infusion of solution must be completed within 24 hours of preparation. The ready-to-use bottles are for single use only; if any preparation is necessary (preparing reduced dosage for patients with renal impairment), the prepared, diluted solution may be refrigerated at 2°C to 8°C if not used immediately. Infusion of solution must be completed within 24 hours of preparation. The previously withdrawn volume from the ready-to-use solution should be discarded; do not store or reuse.	No	Yes
Zoledronic acid (Reclast)	Osteoporosis	IV	No	No	Yes	Store at room temperature of 25°C; excursions permitted to 15°C to 30°C. After opening, stable for 24 hours at 2°C to 8°C. If refrigerated, allow the refrigerated solution to reach room temperature before administration. Store concentrate vials and ready-to-use bottles at 25°C; excursions permitted to 15°C to 30°C. Diluted solutions for infusion in D5W or NS which are not used immediately after preparation should be refrigerated at 2°C to 8°C. Infusion of solution must be completed within 24 hours of preparation. The ready-to-use bottles are for single use only; if any preparation is necessary (preparing reduced dosage for patients with renal impairment), the prepared, diluted solution may be refrigerated at 2°C to 8°C if not used immediately. Infusion of solution must be completed within 24 hours of preparation. The previously withdrawn volume from the ready-to-use solution should be discarded; do not store or reuse.	No	Yes
Zoledronic acid (Zometa)	Oncology- Injectable	IV	No	No	Yes	reuse.	No	Yes
Zolinza	Oncology- Oral/Topical	Oral	Yes	No	Yes	none	No	Yes
Zomacton	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	(see Tev-Tropin)	No	No

Drug	Treatment For	Dosage Form	Covered as Rx	Self-Injectable Product	Costs \$600 or more for up to a 30 day supply	Special Handling (Refrigerate, Limited Distribution, etc)	Limited Distribution by Manufacturer	Available from Wholesaler
Zorbive	Growth Hormone & Related Disorders	Injection SubQ	Yes	Yes	Yes	Store intact vials and diluent at 15°C to 30°C (59°F to 86°F). Following reconstitution with the provided diluent (bacteriostatic water for injection containing benzyl alcohol), may store under refrigeration at 2°C to 8°C (36°F to 46°F) for up to 14 days; do Not freeze. If reconstituted with sterile water for injection, solution should be used immediately.	No	No
Zydelig	Oncology- Oral/Topical	Oral	Yes	No	Yes	Limited Distribution product Not distributed by CVS Specialty. Dispense in the original container.	Yes	No
Zykadia Zytiga	Oncology- Oral/Topical Oncology- Oral/Topical	Oral Oral	Yes Yes	No No	Yes Yes	Limited Distribution product distributed by CVS Specialty none	Yes No	No Yes